

AMENDMENTS TO THE CLAIMS:

Without prejudice, this listing of the claims replaces all prior versions and listings of the claims in the present application:

LISTING OF CLAIMS:

Claim 1. (Canceled).

2. (Currently Amended) A method for establishing a common key k between a central station Z and a group of subscribers $T1-Tn$, comprising:

providing a publicly known mathematical group G and an element $g \in G$ of a high order in the group G , ~~so that for the group G and the element g a calculation of a discrete logarithm is essentially impossible;~~

using a predetermined threshold method, wherein a random number i is generated by each subscriber Ti of the group of subscribers $T1-Tn$, and from the element $g \in G$ and the random number i , the value g^i is calculated by each subscriber Ti of the group of subscribers $T1-Tn$ and transmitted to the central station Z ; in the central station Z , a random number z is generated; from the random number z and the values g^i , the values $(g^i)^z$ in the group G are calculated, from the values $(g^i)^z$, n shares (s_1, \dots, s_n) of the threshold method are derived, and from the shares (s_1, \dots, s_n) , an $(n, 2n-1)$ threshold method is constructed, a secret of the $(n, 2n-1)$ threshold method being the key k to be established; in the central station Z , $n-1$ further shares $(s_{n+1}, \dots, s_{2n-1})$ differing from shares (s_1, \dots, s_n) are calculated together with the value g^z in the group G and are transmitted to the group of subscribers $T1-Tn$; and for each subscriber Ti of the group of subscribers $T1-Tn$, the key k to be established is reconstructed so that from the value g^z transmitted by the central station Z and the random number i of each subscriber Ti of the group of subscribers $T1-Tn$, the value $(g^z)^i$ in the group G is calculated, and that from the resulting value, applying the $(n, 2n-1)$ threshold method, the share s_i is derived, and that using the share s_i and the further shares $(s_{n+1}, \dots, s_{2n-1})$ transmitted by the central station Z , the key k is reconstructed.

3. (New) The method of claim 2, wherein the group G and the element g are of such large proportions that a calculation of a discrete logarithm is virtually impossible.